

Literatur Kapitel 6 - Schwindel

1. Obholz H-H, Jendyk R. DEGAM-S3-Leitlinie Nr. 17 - Akuter Schwindel (Kurzfassung). 2015.

6.1. Peripher-vestibulärer Schwindel

1. Nimensivu R. EbM-Guidelines: Benigner paroxysmaler Lagerungsschwindel [Internet]. 2017. Verfügbar unter: <https://www.ebm-guidelines.com/ebmga/home> [Stand: 08.08.2018].
2. Leitlinien für Schwindel -Therapie in der Neurologie. Deutsche Gesellschaft für Neurologie. 2015;(September 2012):1–15.
3. Wiest G, Deecke L. Der benigne paroxysmale Lagerungsschwindel. *Journal für Neurologie, Neurochirurgie und Psychiatrie.* 2004;5(3):24–9.
4. Sekitani T, Imate Y, Noguchi T, Inokuma T. Vestibular neuronitis: Epidemiological survey by questionnaire in Japan. *Acta Otolaryngologica.* 1993;113(S503):9–12.
5. Adamec I, Krbot Skoric M, Handzic J, Habek M. Incidence, seasonality and comorbidity in vestibular neuritis. *Neurol Sci* 2015; 36(1):91–5. doi: 10.1007/s10072-014-1912-4.
6. Strupp M, Kremmyda O, Bremova T, Teufel J. Aktuelles zur Pharmakotherapie von Schwindel und Nystagmus. Bd. 31, *Arzneimitteltherapie.* 2013. S. 147–55.
7. Strupp M, Brandt T. Diagnosis and treatment of vertigo and dizziness. *Deutsches Arzteblatt international.* 2008;105(10):173–80.
8. Koors PD, Thacker LR, Coelho DH. Investigation of seasonal variability of vestibular neuronitis. *The Journal of laryngology and otology.* 2013;127(10):968–71.
9. Jeong S-H, Kim H-J, Kim J-S. Vestibular neuritis. *Seminars in neurology.* 2013;33(3):185–94: Arbusow V, Schulz P, Strupp M, Dieterich M, Von Reinhardstoettner A, Rauch E, u. a. Distribution of herpes simplex virus type 1 in human geniculate and vestibular ganglia: Implications for vestibular neuritis. *Annals of Neurology.* 1999;46(3):416–9.
10. Strupp M. Leitlinien für Schwindel-Diagnostik in der Neurologie. Deutsche Gesellschaft für Neurologie. 2012; *Hirnnerven:*1–17.
11. Strupp M, Arbusow V, Maag KP, Gall C, Brandt T. Vestibular exercises improve central vestibulospinal compensation after vestibular neuritis. *Neurology.* 1998;51(3):838–44.
12. McDonnell MN, Hillier SL. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst Rev.* 2015;36(3):248–9.
13. Adrion C, Fischer CS, Wagner J, Gürkov R, Mansmann U, Strupp M. Efficacy and safety of betahistine treatment in patients with Ménière's disease: Primary results of a long term, multicentre, double blind, randomised, placebo controlled, dose defining trial (BEMED trial). *BMJ (Online).* 2016;352.
14. Thirlwall AS, Kundu S. Diuretics for Ménière's disease or syndrome. *Cochrane Database of Systematic Reviews (Online).* 2006;3:CD003599.
15. Nevoux J, Barbara M, Dornhoffer J, Gibson W, Kitahara T, Darrouzet V. International consensus (ICON) on treatment of Ménière's disease. *European Annals of Otorhinolaryngology, Head and Neck Diseases.* 2018.
16. Flores García M, Llata Segura C, Cisneros Lesser J, Pane Pianese C. Endolymphatic Sac Surgery for Ménière's Disease – Current Opinion and Literature Review. *International Archives of Otorhinolaryngology.* 2017;21(2):179–83.
17. Jouko Kotimäki. Morbus Meniere [Internet]. EbM Guidelines. 2017. [Stand: 08.08.2018]. Verfügbar unter: <https://www.ebm-guidelines.com/ebmga/home>.
18. Garduño-Anaya MA, De Toledo HC, Hinojosa-González R, Pane-Pianese C, Ríos-Castañeda LC. Dexamethasone inner ear perfusion by intratympanic injection in unilateral Ménière's disease: A two-year prospective, placebo-controlled, double-blind, randomized trial. *Otolaryngology - Head and Neck Surgery.* 2005;133(2):285–94.
19. Casani A Pietro, Piaggi P, Cerchiai N, Seccia V, Franceschini SS, Dallan I. Intratympanic treatment of intractable unilateral ménière disease: Gentamicin or dexamethasone? A randomized controlled trial. *Otolaryngology - Head and Neck Surgery.* 2012;146(3):430–7.
20. Ul Shamas I. Short term results of intra tympanic gentamicin and dexamethasone on hearing and tinnitus in meniere's disease: A case control study. In: *International Tinnitus Journal.* 2017. S. 21–3.
21. Postema RJ, Kingma CM, Wit HP, Albers FWJ, Van Der Laan BF a M. Intratympanic gentamicin therapy for control of vertigo in unilateral Menire's disease: a prospective, double-blind, randomized, placebo-controlled trial. *Acta oto-laryngologica.* 2008;128(8):876–80.

22. Stokroos R, Kingma H. Selective vestibular ablation by intratympanic gentamicin in patients with unilateral active Ménière's disease: A prospective, double-blind, placebo-controlled, randomized clinical trial. *Acta Oto-Laryngologica*. 2004;124(2):172–5.
23. Pullens B, van Benthem PP. Intratympanic gentamicin for Ménière's disease or syndrome. *Cochrane database of systematic reviews (Online)*. 2011;(3):CD008234.
24. Flanagan S, Mukherjee P, Tonkin J. Outcomes in the use of intra-tympanic gentamicin in the treatment of Meniere's disease. *J Laryngol Otol*. 2006;120(2):98–102.
25. Tarnutzer AA, Bockisch C, Buffone E, Weiler S, Bachmann LM, Weber KP. Disease-specific sparing of the anterior semicircular canals in bilateral vestibulopathy. *Clinical Neurophysiology*. 2016;127(8):2791–801.
26. Zingler VC, Weintz E, Jahn K, Huppert D, Cnyrim C, Brandt T, u. a. Causative factors, epidemiology, and follow-up of bilateral vestibulopathy. In: *Annals of the New York Academy of Sciences*. 2009. S. 505–8.
27. Strupp M, Kim JS, Murofushi T, Straumann D, Jen JC, Rosengren SM, u. a. Bilateral vestibulopathy: Diagnostic criteria consensus document of the classification committee of the barany society. *Journal of Vestibular Research: Equilibrium and Orientation*. 2017;27(4):177–89.
28. Zingler VC, Weintz E, Jahn K, Mike A, Huppert D, Rettinger N, u. a. Follow-up of vestibular function in bilateral vestibulopathy. *Journal of Neurology, Neurosurgery and Psychiatry*. 2008;79(3):284–8.
29. Brandt T, Strupp M, Dieterich M. Vestibular paroxysmia: a treatable neurovascular cross-compression syndrome. Bd. 263, *Journal of Neurology*. 2016. S. 90–6.
30. Hüfner K, Barresi D, Glaser M, Linn J, Adrion C, Mansmann U, u. a. Vestibular paroxysmia: Diagnostic features and medical treatment. *Neurology*. 2008;71(13):1006–14.
31. Strupp M, Dieterich M, Brandt T, Feil K. Therapy of Vestibular Paroxysmia, Superior Oblique Myokymia, and Ocular Neuro-myotonia. Bd. 18, *Current Treatment Options in Neurology*. 2016.

6.2. Zentral-vestibulärer Schwindel

1. Pinna MH, Bento RF, Neto RVdB. Vestibular schwannoma: 825 cases from a 25-year experience. *Int Arch Otorhinolaryngol* 2012; 16(4):466–75. doi: 10.7162/S1809-97772012000400007.
2. Propp JM, McCarthy BJ, Davis FG, Preston-Martin S. Descriptive epidemiology of vestibular schwannomas. *Neuro-oncol* 2006; 8(1):1–11. doi: 10.1215/S1522851704001097.
3. Muzevic D, Legcevic J, Splavski B, Cayé-Thomasen P. Stereotactic radiotherapy for vestibular schwannoma. *Cochrane Database Syst Rev* 2014; (12):CD009897. doi: 10.1002/14651858.CD009897.pub2.
4. Stayman A, Nogueira RG, Gupta R. Diagnosis and management of vertebrobasilar insufficiency. *Curr Treat Options Cardiovasc Med* 2013; 15(2):240–51. doi: 10.1007/s11936-013-0228-7.
5. Savitz SI, Caplan LR. Vertebrobasilar disease. *N Engl J Med* 2005; 352(25):2618–26. doi: 10.1056/NEJMra041544.
6. Bogousslavsky J, van Melle G, Regli F. The Lausanne Stroke Registry: analysis of 1,000 consecutive patients with first stroke. *Stroke* 1988; 19(9):1083–92.
7. Moulin T, Tatu L, Vuillier F, Berger E, Chavot D, Rumbach L. Role of a stroke data bank in evaluating cerebral infarction subtypes: patterns and outcome of 1,776 consecutive patients from the Besançon stroke registry. *Cerebrovasc Dis* 2000; 10(4):261–71. doi: 10.1159/000016068.
8. Moubayed SP, Saliba I. Vertebrobasilar insufficiency presenting as isolated positional vertigo or dizziness: a double-blind retrospective cohort study. *Laryngoscope* 2009; 119(10):2071–6. doi: 10.1002/lary.20597.
9. Kim GW, Heo JH. Vertigo of cerebrovascular origin proven by CT scan or MRI: pitfalls in clinical differentiation from vertigo of aural origin. *Yonsei Med J* 1996; 37(1):47–51. doi: 10.3349/yjmj.1996.37.1.47.

6.3. Somatoformer Schwindel

1. Dieterich M, Staab JP. Functional dizziness: from phobic postural vertigo and chronic subjective dizziness to persistent postural-perceptual dizziness. *Curr Opin Neurol* 2017; 30(1):107–13. doi: 10.1097/WCO.0000000000000417.
2. Eckhardt-Henn A, Tschan R, Best C, Dieterich M. Somatoforme Schwindelsyndrome. *Nervenarzt* 2009; 80(8):909–17. doi: 10.1007/s00115-009-2736-y.
3. Yardley L, Luxon LM, Haacke NP. A longitudinal study of symptoms, anxiety and subjective well-being in patients with vertigo. *Clin Otolaryngol Allied Sci* 1994; 19(2):109–16.

4. Jacob RG. Panic disorder and the vestibular system. *Psychiatr Clin North Am* 1988; 11(2):361–74.
5. Schaaf H. Psychogener Schwindel in der HNO-Heilkunde. Bd. 49, HNO. 2001. S. 307–15.
6. Ardiç FN, Ateşci FC. Is psychogenic dizziness the exact diagnosis? *Eur Arch Otorhinolaryngol* 2006; 263(6):578–81. doi: 10.1007/s00405-006-0013-2.

6.4. Weitere Schwindelformen

1. Fernández L, Breinbauer HA, Delano PH. Vertigo and Dizziness in the Elderly. *Front Neurol* 2015; 6:144. doi: 10.3389/fneur.2015.00144.
2. Cabrera Kang CM, Tusa RJ. Vestibular rehabilitation: rationale and indications. *Semin Neurol* 2013; 33(3):276–85. doi: 10.1055/s-0033-1354593.
3. Howe TE, Rochester L, Neil F, Skelton DA, Ballinger C. Exercise for improving balance in older people. *Cochrane Database Syst Rev* 2011; (11):CD004963. doi: 10.1002/14651858.CD004963.pub3.
4. Susanto M. Dizziness: If not vertigo could it be cardiac disease? *Australian Family Physician*. 2014;43(5):264–9.
5. Kang MH, Xu Y, Wang C, Wu LJ, Lin P, Li F, u. a. Causes of unexplained syncope in children. *Chinese Journal of Contemporary Pediatrics*. 2012;14(10):771–4.
6. Brignole M, Menozzi C, Bartoletti A, Giada F, Lagi A, Ungar A et al. A new management of syncope: prospective systematic guideline-based evaluation of patients referred urgently to general hospitals. *Eur Heart J* 2006; 27(1):76–82. doi: 10.1093/eurheartj/ehi647.
7. ECTOR H, REYBROUCK T, HEIDBUCHEL H, GEWILLIG M, WERF F. Tilt Training: A New Treatment for Recurrent Neurocardiogenic Syncope and Severe Orthostatic Intolerance. *Pacing Clin Electro* 1998; 21(1):193–6. doi: 10.1111/j.1540-8159.1998.tb01087.x.
8. Hülse M. Der zervikogene Schwindel. In: W. S, Herausgeber. *Schwindel und schwindelbegleitende Symptome*. Vienna: Springer; 1994.
9. Li Y, Peng B. Pathogenesis, Diagnosis, and Treatment of Cervical Vertigo. *Pain Physician* 2015; 18(4):E583–95.
10. Yacovino DA, Hain TC. Clinical characteristics of cervicogenic-related dizziness and vertigo. *Semin Neurol* 2013; 33(3):244–55. doi: 10.1055/s-0033-1354592.
11. Wrisley DM, Sparto PJ, Whitney SL, Furman JM. Cervicogenic dizziness: a review of diagnosis and treatment. *J Orthop Sports Phys Ther* 2000; 30(12):755–66. doi: 10.2519/jospt.2000.30.12.755.
12. Karlberg M, Magnusson M, Malmström EM, Melander A, Moritz U. Postural and symptomatic improvement after physiotherapy in patients with dizziness of suspected cervical origin. *Arch Phys Med Rehabil* 1996; 77(9):874–82.
13. Reid SA, Rivett DA. Manual therapy treatment of cervicogenic dizziness: a systematic review. *Man Ther* 2005; 10(1):4–13. doi: 10.1016/j.math.2004.03.006.